



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification<sup>4</sup> :</b>  <b>C07D 209/18, A61K 31/405</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 88/ 09785</b>  <b>(43) International Publication Date:</b> <b>15 December 1988 (15.12.88)</b>
<b>(21) International Application Number:</b> PCT/IT88/00041 <b>(22) International Filing Date:</b> 1 June 1988 (01.06.88)  <b>(31) Priority Application Number:</b> 48014 A/87 <b>(32) Priority Date:</b> 3 June 1987 (03.06.87) <b>(33) Priority Country:</b> IT  <b>(71) Applicant (for all designated States except US):</b> POLI-FARMA S.P.A. [IT/IT]; Via Tor Sapienza, 138, I-00155 Roma (IT). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only) :</b> DE LUCA, Giovanna [IT/IT]; Via Ugo de Carolis, 124, I-00136 Roma (IT). DI STAZIO, Giovanni [IT/IT]; Via Clivo di Cinnna, 221, I-00136 Roma (IT). MARGONELLI, Andrea [IT/IT]; Via G. Massaia, 65, I-00154 Roma (IT). MATERAZZI, Mario [IT/IT]; Via Delleani, 10, I-00155 Roma (IT). POLITI, Vincenzo [IT/IT]; Via Albano, 77, I-00179 Roma (IT).		<b>(74) Agents:</b> BAZZICHELLI, Alfredo et al.; Società Italiana Brevetti S.p.A., Piazza Poli, 42, I-00187 Roma (IT).  <b>(81) Designated States:</b> AT (European patent), AU, BE (European patent), BR, CH (European patent), DE (European patent), DK, FI, FR (European patent), GE (European patent), IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent), US.  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> 3-INDOLEPYRUVIC ACID DERIVATIVES AND PHARMACEUTICAL USE THEREOF		
<b>(57) Abstract</b>  3-indolepyruvic acid and derivatives thereof in the form of salts, esters and amides are compounds pharmaceutically active as agents for antagonising disturbances of the central nervous system, such as epilepsy, cerebral ischemia, ictus and Alzheimer's disease, through a mechanism of action which raises the level of kynurenic acid in the brain.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

- 1 -

**"3-INDOLEPYRUVIC ACID DERIVATIVES AND PHARMACEUTICAL  
USE THEREOF"**

&gt;&lt;&gt;&lt;&gt;&lt;&gt;&lt;

**DESCRIPTION**

The present invention relates to novel compounds which are derivatives, in the form of salts, esters and amides, of 3-indolepyruvic acid.

5 The present invention further relates to the use of 3-indolepyruvic acid, and the above mentioned derivatives thereof, as pharmaceutically active agents for the treatment of disturbances of the central nervous system produced in the brain by the presence of a great  
10 amount of superoxide anions or free radicals. Representatives of such disturbances are epilepsy, cerebral ischemia, ictus and Alzheimer's disease.

It has been shown that a pharmaceutical administration of the compounds according to the invention produces  
15 at the brain level an increase of kynurenic acid, which has been recognised as a natural antagonist of the excitatory aminoacids.

References to the prior art

3-indolepyruvic acid is a known compound.

20 In published European Patent Application N° 106813, corresponding to USA Patent N° 4551471, a process for the enzymatic synthesis of 3-indolepyruvic acid is described, which uses aspartate aminotransferase as an enzyme.

25 In published European Application N° 0227787 a process for the chemical synthesis of 3-indolepyruvic acid is described, which uses a coupling reaction starting from L-tryptophan.

Moreover, it has been known for several years that  
30 tryptophan derivatives can interact in mammals with the receptors of so-called "excitatory aminoacids", namely glutamate, N-methyl-aspartate, ibotenic acid,

- 2 -

kainic acid, and so on.

5 In particular it has been observed that quinolinic acid and kynurenine are powerful agonists of excitatory aminoacids present in the central nervous system of mammals (as an example see: Science 219, 316-8, 1983; Neuropharmacology 23, 333-7, 1984) and they can lead to a rapid neuronal decay similar to that observed in epilepsy and Huntington's Chorea (TIPS 1984, page 215).

10 The clinical importance of the excitatory aminoacids has become of considerable interest, such as to induce the organization of periodical symposiums for up-dating, such as the one held in London on 13th and 14th April 1987 ("Excitatory aminoacids in health and disease") and the publication of full monographic editions in  
15 scientific magazines (see TINS, Volume 10, N° 7, 1987). According to the most recent views, the inhibitors of excitatory aminoacids can be used for blocking tremor and spasticity, epilepsy, neurodegenerative disorders, cerebral ischemia, psychosis, and the consequences of  
20 cerebral ictus (Scrip 1198, 27, 1987).

It has recently been discovered that a further derivative of tryptophan, namely kynurenic acid, is a powerful inhibitor of the excitatory effects manifested by quinolinic acid, N-methyl-D-aspartate and other aminoacids  
25 (J. Pharmacol. Exptl. Ther. 236, 293-9, 1986) and it could therefore hinder the neuronal decay in the brain produced by the above mentioned compounds during epileptic attacks, ictus, cerebral ischemia and more generally in neuronal alteration effects occurring during ageing.

30 However, it is not possible to obtain an increase of kynurenic acid at the brain level by means of an administration of kynurenic acid itself, in that the blood-brain barrier prevents the compound from  
35 reaching the brain from the outside. Consequently it cannot be usefully administered either orally or via

- 3 -

normal injection, to increase the cerebral level of kynurenic acid.

It is known, moreover, from the prior art that kynurenic acid is an endogenous compound. It is originated in very small amounts from food tryptophan. However, not even tryptophan can be administered as a drug for this use, in that it is also transformed into kynurenines which have an action which antagonizes that of kynurenic acid.

It has been shown previously (published European Application N° 106813) that 3-indolepyruvic acid is a privileged precursor of cerebral serotonin and, as such, it could be used pharmacologically in all the diseases characterized by a lack of such endogenous amine.

#### Summary of the invention

It has now been surprisingly found that 3-indolepyruvic acid, as well as many simple derivative thereof, can be easily transformed "in vivo" in mammals into kynurenic acid and that the cerebral level of this acid can be substantially increased.


In fact it is surprising and it is an extremely useful result, in accordance with the hereinbefore exposed considerations, that the level of kynurenic acid in the brain can be increased by administration of 3-indolepyruvic acid or a compound according to the invention.

On the basis of pharmacological tests which will be hereinafter reported, which evidence an increase of kynurenic acid at the brain level following an administration of the compounds according to the invention, an explanation is given hereinbelow of the possible mechanism of action.

The fact that 3-indolepyruvic acid is easily transformed into kynurenic acid in all the organs subjected to our tests, whereas their derivatives of the kynurenine

- 4 -

class are absent, suggests that the ket acid undergoes the following transformations in the mammal organism:



The opening of the indole ring of 3-indolepyruvic acid could be produced through indoleamine-dioxygenase enzyme, which is known to be present in all mammal tissues and to act only in the presence of superoxide anions (see for example: J.Biol.Chem. 250, 5960-6, 1975). However, a direct opening of the ring even in the absence of enzymes cannot be excluded, as it is known that the superoxide anions and the free radicals are extremely active compounds and the formation of kynurenic acid from 3-indolepyruvic acid has also been ascertained in "in vitro" systems without any presence of mammal tissues (see Test N° 1 hereinafter reported).

15

20

As it is generally proved that the superoxide anions and the free radicals are involved in the raising of a plurality of situations of neuronal decay (see "Free radicals in medicine and biology", Acta Physiol. Scand. suppl. 492, 1980), 3-indolepyruvic acid and the derivatives thereof appear to be selective agents for increasing the levels of kynurenic acid in the brain, mainly in situations of cerebral disturbance, and consequently they can be used as drugs in pathologies such as epilepsy, ictus, cerebral ischemia and dementia senilis of Alzheimer's type.

25

30


A process for the production of 3-indolepyruvic acid can be effected starting from L-tryptophan, comprising

35

- 5 -

reacting L-tryptophan with L-aminoacid oxydas enzyme in an oxygen containing aqueous medium at a pH of about 7,5.

Object of the invention are novel derivatives of 3-indolepyruvic acid of formula



in which X is -OR, -NHR, -NR<sub>2</sub>, or  

$$\begin{array}{c} \text{-NH-CH-COO-} \\ | \\ \text{R}' \end{array}$$

wherein R is methyl, ethyl, propyl, isopropyl, butyl, tert.butyl, cyclohexyl, or benzyl,

R' is a group which, together with the group -NH-CH-COO- forms an aminoacid radical,

and R'' is H, or methyl, ethyl, propyl, isopropyl, butyl, tert.butyl, cyclohexyl, or benzyl,

and the alkali and earth-alkali metal salts of 3-indolepyruvic acid.

A further object of the invention are pharmaceutical formulations containing the above mentioned novel compounds for the treatment of disturbances of the central nervous system produced by the presence in the brain of a large amount of superoxide anions or free radicals, and the use of 3-indolepyruvic acid and the derivatives thereof of formula 1, for the manufacture of medicaments for the above mentioned therapeutical indication.

#### Description of the process for the production of 3-indolepyruvic acid

3-indolepyruvic acid can be prepared starting from L-tryptophane, taking advantage of a new method of enzymatic synthesis. In fact the amino acid is reacted

- 6 -

with L-aminoacid oxydase, an enzyme which is present in a large amount in a number of snake venoms, in a continuously oxygenated aqueous medium.

3-indolepyruvic acid is formed according to the following reaction:



To avoid the degradation by  $\text{H}_2\text{O}_2$  of the ketoacid formed, catalase, an enzyme which transforms  $\text{H}_2\text{O}_2$  into  $\text{H}_2\text{O}$  and  $\text{O}_2$ , is also added to the solution.

A considerable amount of 3-indolepyruvic acid can be obtained in a continuous flow reactor comprising a first column filled with resin to which a suitable amount of L-aminoacid oxydase and catalase enzymes have been linked.

L-aminoacid oxydase enzyme can be obtained from the venom of several snake species, such as: Ankistrodon, Bitis, Crotalus, Dendroaspis, Naja, Pseudechis, Trimeresurus, viper, and the like.

Catalase can be obtained from cattle liver, for example.

It has been shown that a plurality of resins are able to link the enzymes, in such a way as not to affect their catalytic activity. Among them, particularly useful are Eupergit C type resin (produced by Rohm Pharma), Aminoaril CPG (produced by Pearce), Michroprep silica (produced by E. Merck) and sodium alginate coated with quaternized polyethylene-imine.

In the second column an ion exchange resin is charged (such as Amberlite X AD-4), able to hold 3-indolepyruvic acid formed and allow the passage of tryptophan to be recycled.

The system was activated by two peristaltic pumps of the same flow rate, connected to the first column and to the reactant supply, consisting of a buffered L-tryptophan solution (such as phosphate buffer with

- 7 -

a pH from 7 to 8).

Hereinafter an example of preparation of 3-indolepyruvic acid is described.

500 mg L-aminoacid oxydase enzyme from venom of  
5 Crotalus Atrox were dissolved into phosphate buffer  
1M pH 7,5.

6 g Eupergit C resin were added under light stirring  
and left standing for 24 hours at room temperature.  
At the same time 10 mg of catalase from cattle liver  
10 were added to 10 ml of 1M phosphate buffer pH 7,5.

3 g Eupergit C resin were then added under light  
stirring and left standing for 24 hours at room temperature.  
The resins were united in a porous filter and  
the solution was eluted by gravity. After washing with  
15 300 ml of 0,1 M phosphate buffer pH 7,5, the resin was  
packed into a first column. A second column was charged  
with 5 g Amberlite X AD-4 resin, previously adjusted  
with 0,1 M phosphate buffer pH 7,5. 500 mg L-tryptophan  
in 100 ml of 0,1 M phosphate buffer pH 7,5 were added  
20 in the solution supply. The flow rate was set at 1  
ml/minute. After 22 hours the second column was cut  
off and the solution was eluted with methanol.

The 3-indolepyruvic acid thus obtained was dried  
and weighed (about 200 mg).

25 Preparation of the derivatives of 3-indolepyruvic  
acid

The 3-indolepyruvic acid derivatives were synthesised  
with the aim of enabling the compounds to pass more  
easily through the blood-brain barrier and consequently  
30 to increase the levels of kynurenic acid formed  
in the central nervous system.

For this synthesis four routes were essentially  
followed:

a) the salts were obtained by the alcoholates of  
35 the corresponding metals in anhydrous ether;

- 8 -

b) methyl ester of 3-indolepyruvic acid was produced by reacting the acid with diazomethane;

c) methyl ester and other esters were produced by reacting the carboxylic ion with alkyl halides, with corresponding nucleophilic substitution of the halogen. As the acid is highly unstable, a salification reaction was initially effected with DBU (1,8-diazabicyclo-[5.4.0]-undec-7-ene), which forms a complex with a large charge displacement. This complex, which is strongly nucleophilic and a weak base, can attack primary, secondary, tertiary alkyl and benzyl halides, without giving rise to simultaneous elimination reactions. In this manner methyl, ethyl, isopropyl, tert.butyl, benzyl esters were synthesized and in a similar manner propyl, butyl, cyclohexyl esters and the like can be synthesized;

d) amide type compounds were obtained by a reaction activating the carboxyl group of 3-indolepyruvic acid by means of a carbodiimide, in the presence of hydroxybenzotriazole, with formation of the corresponding activated ester. This is attacked by an amine group with formation of the corresponding amide bond (when an amine was dealt with) or a peptide bond (when a protected amino acid was dealt with).


Hereinafter some examples of preparation for the 3-indolepyruvic acid derivatives are described.

#### Example 1

Preparation of the Mg salt of 3-indolepyruvic acid.

The compound was prepared by reacting the ketoacid with magnesium oxide in an anhydrous ether medium:

30



An anhydrous ether solution of 3-indolepyruvic acid (IPA) (200 mg) is added with 40 mg MgO under stirring and argon environment, at 20-25°C. The suspension was

35

- 9 -

stirred for on night. A large amount of flocculent water soluble precipitate was recovered, which was carefully washed with Et<sub>2</sub>O. The precipitate was dried under vacuum for one night. The conversion into salt was about 90%. This salt shows a 215°C melting point with decomposition.

#### Example 2

Sodium, potassium, calcium and barium 3-indolepyruvates are produced in the same manner as described in Example 1.

#### Example 3

Preparation of 3-indolepyruvic acid methyl ester.

1,0 g of 3-indolepyruvic acid were suspended into 50 ml anhydrous benzene under argon atmosphere.

1,5 ml DBU were added at room temperature, under brisk stirring. Heating was then started for refluxing benzene.

Almost immediately the yellow suspension changes into a dark yellow rubbery solid, which makes stirring difficult. At a temperature of about 50 to 60°C 1,0 ml CH<sub>3</sub>I diluted in 5 ml anhydrous benzene were added. On refluxing benzene, it was noted that the rubbery solid was transformed into a brown-red compound.

After one night refluxing under argon atmosphere, the benzene solution was decanted and the reddish solid was taken with CH<sub>2</sub>Cl<sub>2</sub> until dissolved. The organic solutions were united and the precipitate was filtered off. The organic solution was washed successively with 5% solution NaHCO<sub>3</sub> (10 x 50 ml), saturated NaCl solution up to neutrality, 5% solution of KHSO<sub>4</sub> (10 x 50 ml) and saturated NaCl solution up to neutrality. After leaving for one night on anhydrous Na<sub>2</sub>SO<sub>4</sub>, this was filtered off and the residue was purified by flash chromatography on silica gel (eluent CH<sub>2</sub>Cl<sub>2</sub> 90/acetone 10),

- 10 -

thus obtaining a lightly yellow chromatographically pure solid (on spot on TLC and on peak in HPLC).

The structure was confirmed by IR and NMR spectrometry. The yield was about 60%.

5        Example 4

Preparation of 3-indolepyruvic acid ethyl ester.

Following the same method as previously described, 1,0 g 3-indole-pyruvic acid were reacted with 1,5 ml DBU and 790 microliters ethylbromide. In this case  
10 the benzene solution appeared red orange. After flash chromatography purification the ethyl ester was obtained with a yield of about 70%.

Example 5

Preparation of 3-indolepyruvic acid benzyl ester.

15 Following the same method as previously used, 1,0 g 3-indolepyruvic acid were reacted with 1,5 ml DBU and 850 microliters benzylchloride. The benzene solution before the extractions appeared red orange. Flash chromatography purification showed a 65% yield of benzyl ester.

20        Example 6

In a similar manner as described in the preceding examples, the propyl, isopropyl, butyl, tert.butyl and cyclohexyl esters are prepared.

Example 7

25 Preparation of N,N-dimethyl 3-indolepyruvamide.

A solution of 500 mg 3-indolepyruvic acid in 25 ml anhydrous tetrahydrofuran on an ice-salt bath, under argon atmosphere and under stirring was additioned in swift succession with: 400 mg hydroxybenzotriazole (HOBt)  
30 and 570 mg 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide.HCl (EDC.HCl). After dissolving and still maintaining the ice-salt bath, 240 mg dimethylamin .HCl were added dissolved in 10 ml anhydrous tetrahydr furan and 325 microliters 4-methylmorpholine (NMM). The yellowish  
35

- 11 -

gold solution was maintained under stirring for one night, leaving the bath temperature to rise up to room temperature. The volume was then tripled adding  $\text{CH}_2\text{Cl}_2$  and the organic solution was washed in sequence with 5%  $\text{NaHCO}_3$  (10 x 25 ml), saturated  $\text{NaCl}$  solution (3 x 25 ml), 5%  $\text{KHSO}_4$  (10 x 25 ml) and saturated  $\text{NaCl}$  solution up to neutrality. After standing for one night on anhydrous  $\text{Na}_2\text{SO}_4$ , this was filtered off, the mixture was purified by flash chromatography (silica gel, eluent  $\text{CH}_2\text{Cl}_2$  25/n-propanol 1,5).

A strongly hygroscopic solid was obtained which showed itself to be pure on TLC and on HPLC. The amide was characterized by IR and NMR spectra (yield about 35%).

#### Example 8

Preparation of N-cyclohexyl 3-indolepyruvamide.

The same method as described in Example 7 was followed for the preparation of N,N-dimethylamide. 500 mg 3-indolepyruvic acid were reacted with 400 ml HOBt and 570 ml EDC.HCl. 340 microliters of freshly distilled cyclohexylamine and 325 microliters NMM were added to the solution, which was reacted for 5 hours.

After flash chromatography purification, a yield of about 50% was detected.

#### Example 9

Preparation of N-benzyl 3-indolepyruvamide.

Following the same method as previously described in Example 7, 500 mg 3-indolepyruvic acid were reacted with 400 mg HOBt and 570 mg EDC.HCl. 425 mg benzylamine and 325 microliters NMM were then added. After reacting for one night, it was purified by flash chromatography.

The final yield appears to be about 55%.

#### Example 10

Following the same method as described in Examples 7 to 9, N,N-diethyl, N,N-dipropyl, N,N-dibutyl, N-methyl, N-ethyl, N-propyl, N-isopropyl, N-butyl, N-tert.butyl

- 12 -

3-indolepyruvoamide are produced.

Pseudo-dipeptides have been prepared by reaction of 3-indolepyruvic acid (IPA) and methyl esters of amino-acids.

5. Example 11

Preparation of IPA-L-ALA-OMe.

400 mg HOBt and 570 mg EDC.HCl were added in a swift sequence to a solution of 500 mg 3-indolepyruvic acid in 30 ml anhydrous tetrahydrofuran under stirring, on an ice-salt bath and under argon atmosphere. After dissolving and maintaining the ice-salt bath, a cold solution of 415 mg L-alanine methylester.HCl in 10 ml anhydrous tetrahydrofuran and 325 microliters NMM was added. The yellowish solution was stirred for three hours leaving the temperature to rise to room temperature. Then the volume was tripled with CH<sub>2</sub>Cl<sub>2</sub> and the reaction mixture was successively treated with 5% NaHCO<sub>3</sub> (10 x 20 ml), saturated NaCl solution (3 x 50 ml), 5% KHSO<sub>4</sub> (10 x 20 ml) and saturated NaCl solution up to neutrality. The yellowish solution was left standing for one night on anhydrous Na<sub>2</sub>SO<sub>4</sub>, then this was filtered off and the pseudo-dipeptide was purified by flash chromatography.

A hygroscopic yellowish solid was obtained, which appeared pure on TLC and in HPLC. The structure was confirmed by IR and NMR spectra. Yield was 40%.

Following the same method the compound IPA-D-ALA-OMe was prepared, obviously starting from D-alanine.

Both the esters were then unblocked to obtain the free acids.

30. Example 12

Preparation of IPA-L-PHE-OMe, IPA-D-PHE-OMe and the corresponding free acids.

By the same method as previously described, 500 mg 3-indol pyruvic acid were reacted with 400 mg HOBt and 570 mg EDC.HCl. 640 mg phenylalanine methyl ester hydr -

- 13 -

chl ride (L or D) and 325 microliters NMM were then added. After reacting for five hours and purifying, hygroscopic yellowish solids were recovered with a yield of about 55%.

5        Example 13

Preparation of IPA-L-TYR-OMe, IPA-D-TYR-OMe and the corresponding free acids.

By using the same method as previously described, 500 mg of 3-indolepyruvic acid were reacted with 400  
10 mg HOBT and 570 mg EDC.HCl. 690 mg tyrosine methylester (L or D) and 325 microliters NMM were added. After reacting for five hours and purifying, hygroscopic yellowish-gold solids were recovered with a yield of 65%.

Example 14

15        Preparation of IPA-L-PRO-OMe, IPA-D-PRO-OMe and the corresponding free acids.

Following the same method as previously described, 500 mg 3-indolepyruvic acid were reacted with 400 mg HOBT and 570 mg EDC.HCl. 370 mg proline methylester  
20 hydrochloride (L or D) and 270 microliters NMM were added. After reacting for one night and purifying, hygroscopic yellowish solids were recovered with a yield of 40%.

Example 15

25        Following the same method as described in Examples 10 to 14, the other pseudo-dipeptides are obtained by reacting 3-indolepyruvic acid with the corresponding aminoacid methylester.

Pharmacological tests

30        Pharmacological tests have been carried out to evaluate the levels and the activity of kynurenic acid in various animal organs.

For the measurement of such levels after administration of 3-indolepyruvic acid or a derivative thereof, a system  
35 was set up based on ultraviolet detection of the compounds

- 14 -

separated by high pressure liquid chromatography.

Briefly, after sacrifice of the animal, some of the principal organs were taken and homogenized with alkaline ethanol. Proteins and cell remains were removed by centrifugation, whilst the overnatant was adsorbed on Bio-Rad AG 1x8 resin. After washing with water it was eluted with formic acid. It was then passed on Bio-Rad AG 50W-x8 resin, eluted with 3% ammonia, dried and retaken with a small amount of water.

The sample thus obtained was injected into a Waters apparatus for high pressure liquid chromatography, using a C18 10 micron column and eluting with 0.02 M citric acid containing 10% of methanol. The kynurenic acid levels were extrapolated by the calibration line of the acid and reading at 234 nm.

The tests hereinafter reported have the object of illustrating the effect of administration of IPA and the derivatives thereof of formula 1, on the production of kynurenic acid in the brain. The tests also show that the agent producing such an effect is IPA, rather than tryptophan.

It is well known in fact, as mentioned hereinbefore, that tryptophan (briefly TRP) is transformed into kynurenines within the peripheral tissues and a doubt could be raised that these kynurenines could be responsible for the production of kynurenic acid in the brain. This however is not the case, as the following tests show. The tests confirm that only an administration of IPA causes the level of kynurenic acid in the brain to increase substantially.

#### Test 1

Transformation of 3-indolepyruvic acid into kynurenic acid "in vitro", within a non enzymatic system.

When 3-indolepyruvic acid or a derivative thereof of formula 1 is incubated in a slightly acid solution

- 15 -

(HCl 0,1 M), in the presence of oxidising agents ( $O_2$ ) and light, in time the formation of small amounts of kynurenic acid is observed. Particularly starting from 2 mg 3-indolepyruvic acid, 0,5 micrograms are obtained after 2 hours and 0,7 micrograms after 4 hours.

This test shows that 3-indolepyruvic acid is transformed, even without enzymes, into kynurenic acid, even "in vitro".

### Test 2

Transformation of 3-indolepyruvic acid into kynurenic acid in tissue homogenates.

Male rats fasting for 16 hours were sacrificed and their organs were taken and homogenized in 5 volumes of 0,05 M phosphate buffer pH 7,4 + 0,14 M KCl.

Small amounts of homogenate (0,8 ml) were added with various amounts of 3-indolepyruvic acid or the derivatives thereof, dissolved into a buffer solution, and with a system for the production of free radicals, consisting of ascorbic acid (final  $10^{-5}$  M),  $FeCl_3$  (final  $10^{-6}$  M) and  $H_2O_2$  (final  $10^{-4}$  M). It was stirred at  $37^\circ C$ , then the reaction was blocked. The following results were obtained.

TABLE 1

25

## Kidney homogenate

## Amount of kynurenic acid formed

30'      60'      120'

30

3-indolepyruvic acid	final 0.8 mg	160 ng	190 ng	208 ng
3-indolepyruvic acid	final 1.6 mg	232 ng	280 ng	315 ng
3-indolepyruvic acid	final 4 mg	185 ng	221 ng	270 ng
Tryptophan	final 1.6 mg	55 ng	82 ng	111 ng

35

- 16 -

TABLE 2

Brain homogenate

Amount of kynurenic acid formed

		30'	60'	120'
5	3-indolepyruvic acid 1.6 mg	165 ng	280 ng	380 ng
	Tryptophan 1.6 mg	0	0	0
	IPA-Methylester	48 ng	65 ng	83 ng
	IPE-TYR-Ome	44 ng	48 ng	79 ng
	IPE-PHE-Ome	75 ng	82 ng	92 ng

10

IPA = 3-indolepyruvic acid

IPA-TYR-Ome = 3-indolepyruvic acid tyrosine methylester

IPA-PHE-Ome = 3-indolepyruvic acid phenylalanine methylester

15

The results show that kynurenic acid is formed in time when 3-indolepyruvic acid is incubated with homogenates of various organs.

20

Tryptophan in the brain is never transformed into kynurenic acid, as shown in Table 2, and this evidences that the transformation into kynurenic acid only occurs directly from IPA ketoacid.

25

In the kidney, where kynurenine transaminase is present, it is possible that kynurenic acid is also formed from kynurenine. It is known, however, that said kynurenic acid cannot reach the brain, in that this is hindered by the blood-brain barrier.

30

Moreover, Table 2 shows that the derivatives of IPA are also able to produce kynurenic acid when incubated in brain homogenate.

Resuming, this test shows that IPA is able to transform into kynurenic acid in the brain, whereas tryptophan cannot.

Test 3

35

Effect on the content of kynurenic acid after administration of IPA in rat.

- 17 -

Male rats were treated with saline solution (controls) or with various doses of IPA, which was injected 3 hours before sacrifice. Each lot comprised 10 animals.

The results are reported in picomoles/g of tissue (or ml of blood)  $\pm$  standard error.

TABLE 3

	Liver	Blood	Brain
Saline	121 $\pm$ 5	25 $\pm$ 3	18 $\pm$ 2
IPA - 100 mg/Kg i.p.	240 $\pm$ 20	62 $\pm$ 3	24 $\pm$ 1
10 IPA - 100 mg/Kg per os	275 $\pm$ 25	87 $\pm$ 8	28 $\pm$ 2
IPA - 250 mg/Kg i.p.	547 $\pm$ 35	164 $\pm$ 10	32 $\pm$ 3
IPA - 250 mg/Kg per os	485 $\pm$ 40	120 $\pm$ 12	38 $\pm$ 3
IPA - 500 mg/Kg i.p.	2860 $\pm$ 300	442 $\pm$ 20	110 $\pm$ 10

15

The results show that administration of IPA produces an increase in the levels of kynurenic acid in the tissues under test. Such an increase is dose-dependent. This fact is particularly relevant in the brain, where, for a 500 mg/Kg dose of IPA, values of kynurenic acid six times higher than the controls are obtained.

20

This test shows that an administration of IPA causes the level of kynurenic acid in the brain to increase to an extent depending on the dose. It is to be noted that, as previously illustrated, an administration of tryptophan does not produce such an effect.

25

#### Test 4

30

Effects on the content of tryptophan and kynurenic acid in the brain of rats, by treatment with IPA or tryptophan.

Male rats were treated with saline solution (controls), IPA or tryptophan i.p. one hour before sacrifice. Each lot was of six animals.

35

The reported values are in picomoles/g for kynurenic acid and picomoles/mg for tryptophan.

- 18 -

TABLE 4

Kynurenic acid      Tryptophan      KYN/TRPx10<sup>3</sup>

	Saline	18 ± 2	21.3 ± 3	0.86
5	IPA 250 mg.Kg	47 ± 5	124 ± 10	0.40
	TRP 250 mg/Kg	35 ± 3	206 ± 15	0.16

10      The results of this test show that in the brain, kynurenic acid is formed preferably by direct opening of the indole ring of IPA, rather than through the kynurenins.

IPA consequently can be considered as a selective precursor of cerebral kynurenic acid.

15      In fact, as in test N° 2 it was shown that TRP does not transform into kynurenic acid in the brain, the presence of kynurenic acid due to administration of TRP should be due to transformation into kynurenic acid of kynurenines coming from peripheral organs.

20      On the other hand, the ratio of the content of kynurenic acid to tryptophan excludes the same pathway for 3-indolepyruvic acid.

This proved that 3-indolepyruvic acid transforms directly into kynurenic acid in the brain.

#### 25      Test 5

Time course of the kynurenic acid levels after administration of IPA.

Male rats, in lots of five animals each, were treated i.p. with 250 mg/Kg IPA and were sacrificed at different times, evaluating the levels of kynurenic acid present in various organs.

The results are reported in pic moles/g ± standard error.

- 19 -

TABLE 5

	Control	1 hour	2 hours	3 hours	4 hours
Brain	20 $\pm$ 2	39 $\pm$ 4	32 $\pm$ 3	38 $\pm$ 4	28 $\pm$ 3
Liver	80 $\pm$ 10	827 $\pm$ 50	464 $\pm$ 40	235 $\pm$ 5	122 $\pm$ 10
5 Kidney	262 $\pm$ 25	2593 $\pm$ 250	2395 $\pm$ 200	1480 $\pm$ 150	1200 $\pm$ 150
Heart	67 $\pm$ 7	181 $\pm$ 18	177 $\pm$ 20	104 $\pm$ 12	100 $\pm$ 11

The results show a high increase of kynurenic acid in all the organs under test (maximum in kidney). The levels remain high even after 4 hours from administration.

#### Test 6

Effect of administration of IPA or derivatives thereof on audiogenic convulsions in DBA2 mice.

To evaluate whether an increase in kynurenic acid in the brain after administration of IPA could have a protective action on neurons stressed by excitatory aminoacids, the experimental pattern of inducing convulsions in a particular strain of mice (DBA2) by selected sound stimulations was followed. In such pattern kynurenic acid is protective against convulsions.

The substances were injected intravenously one hour before the test. Then the number of animals showing tonic convulsions was measured.

TABLE 6

	Animals with convulsions	%
Controls	9/14	65%
IPA 200 mg/Kg	4/12	33%
Tryptophan 200 mg/Kg	8/10	80%
30 Mg <sup>++</sup> IPA salt	4/14	29%

The results show a good protective action by IPA and its Mg<sup>++</sup> salt against tonic convulsions induced by sound stimulations on DBA2 mice. On the contrary,

- 20 -

tryptophan showed no effective action.

#### Test 7

Action of administration of IPA on NMDA induced convulsions.

5 N-methyl-D-aspartic acid (NMDA) is a compound having a powerful neurotoxic effect, due to activation of specific receptors of the excitatory aminoacids. As it is well known that kynurenic acid protects against the toxicity due to NMDA, the action of IPA, as precursor  
10 of kynurenic acid, was evaluated against the toxicity of such compound.

Mice of Albino Swiss strain were treated with saline (controls) or with IPA (1 g/Kg i.p.) one hour before receiving a toxic dose of NMDA (154 mg/Kg i.p.).

15

TABLE 7

	Tonic convulsions	Deaths
Controls	9/10	8/10
Treated	5/9	3/9

20

From the above results it can be assessed that IPA is able to protect mice against both convulsions and death induced by NMDA.

It follows that 3-indolepyruvic acid and the above  
25 described derivatives thereof, can be used for the treatment of human and animal diseases due to a lack of kynurenic acid at the brain level and more generally for the treatment of disturbances of the central nervous system produced by the excitatory aminoacids.

30 The administration can be made by means of pharmaceutical compositions containing the active substance in a dose of about 2 to 20 mg/Kg body weight for a "per s" administration, and in a dose of about 1 to 10 mg/Kg body weight for a parenteral administration.

35 For the oral, parenteral or rectal administration

- 21 -

the usual pharmaceutical forms can be used, such as pills, capsules, solutions, suspensions, injections, suppositories, in association with pharmaceutically acceptable carriers or diluents and excipients.

5

10

15

20

25


30

35

- 22 -

## CLAIMS

1. 3-indolepyruvic acid derivatives of formula:



in which X is -OR, -NHR, -NR<sub>2</sub>, or




where R is methyl, ethyl, propyl, isopropyl, butyl, tert.butyl, cyclohexyl or benzyl, or a metal selected from the group consisting of K, Mg, Ca, Ba, forming a salt with 3-indolepyruvic acid,

R' is a group which, together with the -NH-CH-COO- group forms an aminoacid radical, and

20 R'' is H, or methyl, ethyl, propyl, isopropyl, butyl, tert.butyl, cyclohexyl or benzyl.

2. Use of a compound of formula



in which X is -OH, -OR, -NHR, -NR<sub>2</sub>, or



where R is methyl, ethyl, propyl, isopropyl, butyl, tert.butyl, cyclohexyl or benzyl, or a metal selected from the group consisting of Na, K, Mg, Ca, Ba, forming a salt with 3-indolepyruvic acid,

R' is a group which, together with the -NH-CH-COO-

- 23 -

group forms an amino acid radical, and

R" is H, or methyl, ethyl, propyl, isopropyl, butyl, tert.butyl, cyclohexyl,


for the manufacture of a medicament for treating  
5 disturbances of the central nervous system due to the presence of excessive amounts of excitatory aminoacids in the brain.

3. Use according to claim 2, in which said disturbances are epilepsy, cerebral ischemia, ictus and Alzheimer's  
10 disease.


4. Use according to claim 2, for the manufacture of a medicament able to increase the cerebral level of kynurenic acid.

5. Pharmaceutical compositions comprising a  
15 3-indolepyruvic acid derivative as defined in formula (1) on claim 1, as an active substance for the treatment of disturbances of the central nervous system due to the presence of an excessive amount of excitatory aminoacids in the brain, in a therapeutically effective amount, and a pharmaceutically acceptable carrier or  
20 diluent.

6. A method for treating disturbances of the central nervous system due to the presence of an excessive amount of excitatory aminoacids in the brain in the mammal,  
25 comprising administering to said mammal a therapeutically effective amount of a compound of formula:



in which X is -OH, -OR, -NHR, -NR<sub>2</sub>, or



- 24 -

where R is methyl, ethyl, propyl, isopropyl, butyl, tert.butyl, cyclohexyl or benzyl, r a metal selected from the group consisting of Na, K, Mg, Ba, forming a salt with 3-indolepyruvic acid,

5 R' is a group which, together with the  $\text{-NH-CH-COO-}$  group forms an aminoacid radical, and

R'' is H, or methyl, ethyl, propyl, isopropyl, butyl, tert.butyl, cyclohexyl or benzyl.

10 7. A method according to claim 6, in which said disturbances are epilepsy, cerebral ischemia, ictus and Alzheimer's disease.

8. A method according to claim 6, for increasing the cerebral level of kynurenic acid.

15

20

25

30

35



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification<sup>4</sup> :</b>  <b>C07D 209/18, A61K 31/405</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 88/09789</b>  <b>(43) International Publication Date:</b> 15 December 1988 (15.12.88)
<b>(21) International Application Number:</b> PCT/IT88/00041 <b>(22) International Filing Date:</b> 1 June 1988 (01.06.88)  <b>(31) Priority Application Number:</b> 48014 A/87 <b>(32) Priority Date:</b> 3 June 1987 (03.06.87) <b>(33) Priority Country:</b> IT  <b>(71) Applicant (for all designated States except US):</b> POLI-FARMA S.P.A. [IT/IT]; Via Tor Sapienza, 138, I-00155 Roma (IT). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only) :</b> DE LUCA, Giovanna [IT/IT]; Via Ugo de Carolis, 124, I-00136 Roma (IT). DI STAZIO, Giovanni [IT/IT]; Via Clivo di Cinnna, 221, I-00136 Roma (IT). MARGONELLI, Andrea [IT/IT]; Via G. Massaia, 65, I-00154 Roma (IT). MATERAZZI, Mario [IT/IT]; Via Delleani, 10, I-00155 Roma (IT). POLITI, Vincenzo [IT/IT]; Via Albano, 77, I-00179 Roma (IT).		<b>(74) Agents:</b> BAZZICHELLI, Alfredo et al.; Società Italiana Brevetti S.p.A., Piazza Poli, 42, I-00187 Roma (IT).  <b>(81) Designated States:</b> AT (European patent), AU, BE (European patent), BR, CH (European patent), DE (European patent), DK, FI, FR (European patent), GB (European patent), IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent), US.  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>  <b>(88) Date of publication of the international search report:</b> 9 February 1989 (09.02.89)
<b>(54) Title:</b> 3-INDOLEPYRUVIC ACID DERIVATIVES AND PHARMACEUTICAL USE THEREOF  <b>(57) Abstract</b>  3-indolepyruvic acid and derivatives thereof in the form of salts, esters and amides are compounds pharmaceutically active as agents for antagonising disturbances of the central nervous system, such as epilepsy, cerebral ischemia, ictus and Alzheimer's disease, through a mechanism of action which raises the level of kynurenic acid in the brain.		

***FOR THE PURPOSES OF INFORMATION ONLY***

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LJ	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

# INTERNATIONAL SEARCH REPORT

International Application No PCT/IT 88/00041

<b>I. CLASSIFICATION F SUBJECT MATTER</b> (If several classification symbols apply, indicate all) <sup>6</sup> According to International Patent Classification (IPC) or to both National Classification and IPC IPC <sup>4</sup> : C 07 D 209/18; A 61 K 31/405																				
<b>II. FIELDS SEARCHED</b> <div style="text-align: center; margin-top: 10px;">Minimum Documentation Searched <sup>7</sup></div> <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 30%; border-bottom: 1px solid black;">Classification System</th> <th style="border-bottom: 1px solid black;">Classification Symbols</th> </tr> <tr> <td style="padding: 5px;">IPC<sup>4</sup></td> <td style="padding: 5px;">C 07 D 209/00</td> </tr> </table> <div style="text-align: center; margin-top: 10px; font-size: small;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>8</sup></div>			Classification System	Classification Symbols	IPC <sup>4</sup>	C 07 D 209/00														
Classification System	Classification Symbols																			
IPC <sup>4</sup>	C 07 D 209/00																			
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT <sup>9</sup></b> <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 10%; border-bottom: 1px solid black;">Category <sup>9</sup></th> <th style="border-bottom: 1px solid black;">Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup></th> <th style="width: 10%; border-bottom: 1px solid black;">Relevant to Claim No. <sup>13</sup></th> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">GB, A, 1550993 (W.R. GRACE &amp; CO.) 22 August 1979 see the whole document --</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1,2</td> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">EP, A, 0048159 (UNIVERSITY OF MIAMI) 24 March 1982 see page 9; page 99, example 706; page 108, compound 43 --</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1,2</td> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">EP, A, 0046291 (MERCK &amp; CO.) 24 February 1982 see pages 1,2; page 58, example 35; page 61, table III --</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1,2</td> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">EP, A, 0046953 (HOECHST AG) 10 March 1982 see page 1,2,9; pages 24-25,41,42 --</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1,2</td> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">Chemical Abstracts, volume 91, no. 3, 16 July 1979, (Columbus, Ohio, US), Yu.V. Erofeev et al.: "Synthesis of esters of 3-indolylpyruvic acid ./.</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1,2</td> </tr> </table>			Category <sup>9</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>	X	GB, A, 1550993 (W.R. GRACE & CO.) 22 August 1979 see the whole document --	1,2	X	EP, A, 0048159 (UNIVERSITY OF MIAMI) 24 March 1982 see page 9; page 99, example 706; page 108, compound 43 --	1,2	X	EP, A, 0046291 (MERCK & CO.) 24 February 1982 see pages 1,2; page 58, example 35; page 61, table III --	1,2	X	EP, A, 0046953 (HOECHST AG) 10 March 1982 see page 1,2,9; pages 24-25,41,42 --	1,2	X	Chemical Abstracts, volume 91, no. 3, 16 July 1979, (Columbus, Ohio, US), Yu.V. Erofeev et al.: "Synthesis of esters of 3-indolylpyruvic acid ./.	1,2
Category <sup>9</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>																		
X	GB, A, 1550993 (W.R. GRACE & CO.) 22 August 1979 see the whole document --	1,2																		
X	EP, A, 0048159 (UNIVERSITY OF MIAMI) 24 March 1982 see page 9; page 99, example 706; page 108, compound 43 --	1,2																		
X	EP, A, 0046291 (MERCK & CO.) 24 February 1982 see pages 1,2; page 58, example 35; page 61, table III --	1,2																		
X	EP, A, 0046953 (HOECHST AG) 10 March 1982 see page 1,2,9; pages 24-25,41,42 --	1,2																		
X	Chemical Abstracts, volume 91, no. 3, 16 July 1979, (Columbus, Ohio, US), Yu.V. Erofeev et al.: "Synthesis of esters of 3-indolylpyruvic acid ./.	1,2																		
<div style="display: flex; justify-content: space-between; font-size: x-small;"> <div style="width: 45%;"> <p><sup>10</sup> Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Z" document member of the same patent family</p> </div> </div>																				
<b>IV. CERTIFICATION</b> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; padding: 5px;">           Date of the Actual Completion of the International Search  <b>4th October 1988</b> </td> <td style="width: 50%; padding: 5px;">           Date of Mailing of this International Search Report  <b>13. 01. 89</b> </td> </tr> <tr> <td style="padding: 5px;">           International Searching Authority  <b>EUROPEAN PATENT OFFICE</b> </td> <td style="padding: 5px;">           Signature of Authorized Officer    <b>P.C.G. VAN DER PUTTEN</b> </td> </tr> </table>			Date of the Actual Completion of the International Search <b>4th October 1988</b>	Date of Mailing of this International Search Report <b>13. 01. 89</b>	International Searching Authority <b>EUROPEAN PATENT OFFICE</b>	Signature of Authorized Officer  <b>P.C.G. VAN DER PUTTEN</b>														
Date of the Actual Completion of the International Search <b>4th October 1988</b>	Date of Mailing of this International Search Report <b>13. 01. 89</b>																			
International Searching Authority <b>EUROPEAN PATENT OFFICE</b>	Signature of Authorized Officer  <b>P.C.G. VAN DER PUTTEN</b>																			

## FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

X	<p>and tryptophan based on <math>\beta</math>-(3-indolyl)-<math>\alpha</math>-nitroacrylic acid esters", see page 708, abstract 21026g, &amp; Zh. Vses. Khim. O-va. 1979, 24(1), 91-2</p> <p>-----</p> <p>Chemical Abstracts, volume 75, no. 13, 27 September 1971, (Columbus, Ohio, US), see page 150, abstract 85992w, &amp; SU, A, 301147 (ALL-UNION SCIENTIFIC-RESEARCH INSTITUTE OF PHYSIOLOGY AND BIOCHEMISTRY OF FARM ANIMALS) 21 April 1971</p> <p>-----</p>	1,2
---	--	-----

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE <sup>1</sup>

This International search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☒ Claim numbers 6-8, because they relate to subject matter not required to be searched by this Authority, namely:

see PCT-Rule 39.1. (iv):

Methods for treatment of the human<sup>2</sup> or animal body by surgery or therapy, as well as diagnostic methods.

2. ☐ Claim numbers \_\_\_\_\_, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claim numbers \_\_\_\_\_, because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING <sup>1</sup>

This International Searching Authority found multiple inventions in this international application as follows:

1. Claims 1 and 2 to 8 where X is not hydrogen.
2. Claims 2 to 4 and 6 to 8 where X is hydrogen

for further information please see PCT/ISA/206 dated 24.10.1988

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.

2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

## Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.  
☒ No protest accompanied the payment of additional search fees.

**ANNEX TO THE INTERNATIONAL SEARCH REPORT  
ON INTERNATIONAL PATENT APPLICATION NO.**

IT 8800041

SA 22652

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 04/01/89. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB-A- 1550993	22-08-79	NL-A- 7710796	11-04-78
		FR-A- 2367045	05-05-78
		DE-A- 2744305	13-04-78
		JP-A- 53046920	27-04-78
		CA-A- 1096881	03-03-81
		CH-A- 624919	31-08-81
		BE-A- 859290	16-01-78
		SE-A- 7711163	08-04-78
EP-A- 0048159	24-03-82	JP-A- 57126468	06-08-82
		AU-A- 7529581	25-03-82
		JP-A- 58035114	01-03-83
EP-A- 0046291	24-02-82	JP-A- 57112373	13-07-82
		US-A- 4680392	14-07-87
		US-A- 4629787	16-12-86
EP-A- 0046953	10-03-82	DE-A- 3032709	29-04-82
		AU-A- 7471881	11-03-82
		AU-B- 544756	13-06-85
		AU-A- 7928487	04-02-88
		EP-A- 0278530	17-08-88
		DE-A- 3118191	25-11-82